



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

PROBLEMS FOR SOLUTION.

ALGEBRA.

333. Proposed by R. D. CARMICHAEL, Princeton University.

Sum the infinite series

$$\frac{1}{(m+1)^2} + \frac{(2m-1)}{(2m+1)^2} + \frac{(3m-1)^2}{(3m+1)^4} + \frac{(4m-1)^3}{(4m+1)^5} + \frac{(5m-1)^4}{(5m+1)^6} + \dots$$

334. Proposed by G. B. M. ZERR, A. M., Ph. D., Philadelphia, Pa.

Sum the series, $2^n - n \cdot 2^{n-2} + \frac{n(n-3)}{2!} 2^{n-4} - \frac{n(n-4)(n-5)}{3!} 2^{n-6}$
 $+ \frac{n(n-5)(n-6)(n-7)}{4!} 2^{n-8} - \frac{n(n-6)(n-7)(n-8)(n-9)}{5!} 2^{n-10} + \dots$

GEOMETRY.

362. Proposed by V. M. SPUNAR, M. and E. E., 3536 Massachusetts Avenue, N. S., Pittsburg, Pa.

Show that the focus of an ellipse may be regarded as an indefinitely small circle having double contact with the ellipse, the directrix being the chord joining the points of contact.

363. Proposed by G. I. HOPKINS, Manchester, N. H.

Construct the triangle, having given, base, vertical angle, and difference between altitude and sum of the other two sides.

364. Proposed by R. C. ARCHIBALD, Providence, R. I.

Between the side of a given rhombus and its adjacent side produced, to insert a straight line of a given length and directed to the opposite corner. [“Euclidean constructions” are particularly desired.]

CALCULUS.

290. Proposed by C. N. SCHMALL, New York City.

When the equation $ax^3 + 2hxy + by^3 + 2gx + 2fy + c = 0$, represents an ellipse, show (by integration) that its area is

$$\frac{\pi (af^2 + bg^2 + ch^2 - abc - 2fgh)}{(ab - h^2)^{\frac{3}{2}}}.$$

291. Proposed by V. M. SPUNAR, M. and E. E., Pittsburg, Pa.

Integrate $\frac{dy}{dx} = ay^2 + bx^m$.

MECHANICS.

244. Proposed by G. B. M. ZERR, A. M., Ph. D., Philadelphia, Pa.

A load P is supported by three strings of equal size lying in the same plane. The middle string is vertical, one string makes with it the angle θ on one side, and the second string makes with it the angle ϕ on the other side. Find the stresses in the strings.